

SEVERE ATTACHMENT DISORDER DEVELOPMENT AND THERAPY FROM EARLY CHILDHOOD TO PUBERTY

1. abstract.
2. family background
3. attachment in the baby.
4. the toddler and pre-school life.
5. school age.
6. puberty.
7. adulthood.
8. late maturation.
9. possible interventions by developmental age.
10. conclusion

By N.P.Rygaard, practicing psychologist, auth. by the Danish Psychologist Association, Denmark

Abstract: The diagnosis of Attachment Disorder (hence: AD) is used when the client is approximately between six and seventeen years of age. However, this “window” is rather narrow, since the causes and symptoms operate very early in life. The author gives an overview of common lifespan events based on clinical experience and studies of AD. An imaginary typical lifespan case, “Henry” is used, and for each stage of development a general description illustrates the typical traits behind the case. The second section gives examples of prevention at each developmental level of the child or youth. The focus is on the period from conception to puberty.

THE DISRUPTION OF HENRY’S SOCIETY

Henry was born into a culture undergoing rapid change. As a consequence, the rituals and roles of caretaking in his family were disrupted by industrialization before he was born: the bonds between his mother and the family were more or less broken up. Where formerly she would have learned the practice of caring for the newborn from her grandmother, her mother and other women, she was more or less on her own during her pregnancy. At school she had been forced to speak English, this also denied her access to the “knowledge of caretaking” which had been accumulated in her native language. She did not let baby Henry sleep in a cradle, but in a bed. She did not breastfeed him, but used a bottle. She did not live in a group of women, carrying Henry in a bag on her back; she had to leave him alone, going to the factory for many hours a day. At the factory she was introduced to alcohol. Henry did not grow up according to tribal tradition; in fact he never met his father or his grandfather.

General aspects: ancient caretaking traditions respond to the baby’s survival challenges in the environment. Whenever the environment changes rapidly the percentage of children having attachment problems tends to rise until new relevant coping traditions for care have been formed. You may study the Inuit, the Danes fifty years ago or any other traditional society transforming rapidly: attachment problems increase whenever the social structure is in the melting pot. For

example, at present the Chinese experience massive migration from rural to urban environments and economies, this produces a host of abandoned children, adopted children and a temporary disruption of the cultural norms of local baby-care patterns and upbringing systems. Our Henry was born in the midst of this turmoil; otherwise he might have been named Atassaq and have had a completely different life. Approximately 3-5 % in a population have signs of severe attachment problems, however in periods of change in society (war, famine, rapid industrialization, migration etc.), the number increases. Most studies in failure of attachment are conducted in the wake of such changes, including John Bowlby's (Bowlby 1969, 1973, 1988) famous studies in attachment with post-war children.

HENRY'S FAMILY

His father was sometimes very kind and sometimes violent towards his mother. He left for good when Henry was six months old. His mother had left home at an early age. She was in many ways still a child, although being 25 when Henry entered the world: she was unable to create a daily rhythm for herself, and emotionally she was unpredictable - she could have fits of rage at one moment and be gay and happy the next.

General aspects: the family pattern producing the AD child is a father who is a) absent, physically or emotionally, b) violent, c) a "serial father" (the mother having many short and superficial relationships with different men). The mother is not particularly young when giving birth; rather she is emotionally unstable and has a fragile personality.

This latter characteristic can be divided into three subgroups: The mother with a disorganized/disorientated attachment pattern (and perhaps many psychopathic traits) (50%), the mother suffering from psychosis during the first years of her child's life (schizophrenia, manic-depressive, borderline and birth psychosis) (40%). Third, the average normal mother (10%) – who either happens to be in crisis due to other events in the first years of the baby's life - or whose baby is so fragile or handicapped at birth that is unable to perceive the mother's attempts to make contact that attachment does not take place in spite of her efforts. Two risk groups in this category are presently growing: adopted children (only if deprived prior to adoption and usually if adopted at age 1 or > age 1), and children with extremely low birth weight (Rygaard 1998).

HENRY IN MOTHER'S WOMB

Henry had odds against him from the start. During pregnancy his mother was overworked and depressed from the loss of her dear ones. Her drinking was toxic to Henry's brain development and she was unable to care for herself. Henry's birth weight was low and the birth itself very complicated. After birth he cried a lot and she did not have much time to comfort him at her breast and have little dialogues with him (Mummy smiles, Henry smiles, Mummy looks gloomy and laughs, Henry looks gloomy and grunts). As soon as he could hold the milk bottle himself, she stopped taking him up when feeding him.

General aspects: most (not all) children who develop AD have experienced what you may call an "insufficient pregnancy and birth". Thus, the combination of a very fragile and neurologically immature newborn can be devastating when combined with subsequent insufficient parenting. Some

children develop AD merely because the parent is unable to be a good caretaker, but in most cases the baby was also fragile from the start. This results in a lack of establishing mutual feedback in the first year of interaction, and this in turn probably results in insufficient neurological growth and in a lack of forming adequate neural networks for emotional and cognitive function and regulation. Genetically, Henry had the misfortune of being a boy, and boys are much more vulnerable to almost any kind of disease and stress than girls. App. 75 % of all children who develop AD will be boys (Schulsinger 1972, McCord 1982).

BABY HENRY

When Baby Henry's mother cuddled him, she often scared him for fun beyond the limits of panic, not being very sensitive to his mood. Sometimes she was loving and kind; sometimes she had sudden fits of rage and yelled at him or shook him, or left him alone if she became engaged in something else. She had a habit of starting to nurse him, and then abruptly put him aside in the midst of all. If he cried or screamed, she would become very excited and scold him for not liking her; as if a baby six months old could have intentions behind his actions. Henry often had stomach problems and colic, he didn't gain weight and height as well as he should, he didn't sleep well and he often had fever. When a health nurse examined him at 9 months, his head circumference was small for age; he was unable to lift his head when lying on the back or stomach, and his muscle tonus was low. She could not make eye contact with him. Consequently he was admitted to a hospital for a month, and a lot of different persons nursed him during the day.

General aspects: Henry's mother displays many behaviour traits of the attachment pattern "Disorganized/ disoriented", typical for mothers who were severely deprived in their own childhood. This attachment pattern is linked to the development of personality disorder later in life. Already at age one there's a 70% chance that Henry will have an identical pattern in adulthood (Shaver & Cassidy 1999).

Neurologically speaking Henry's brain is already far behind:

His brain stem is unable to produce the level of brain activity required because his mother does not touch him or give him vestibular stimulation (which will normally increase brain stem activity and stabilize it at app. 9 months). This means that his nervous system activity remains as unstable and low as in a newborn: pulse, breath, temperature regulation, regulation of sleeping, waking and attention rhythms. Appetite and digestion patterns remain unstable, and this severe condition can quickly become chronic. Also, the brain stem is responsible for growth hormone production, and the lack of stimulation results in a lack of growth (psychosocial dwarfism) and a reduced head circumference.

His "emotional brain" is not being properly programmed. The neurological network that should help him calm down from states of panic and anxiety develops especially from age 10 to 18 months, and the creation of this network depends heavily on the mother's ability to soothe her child and help it regulate the intensity of emotions. Henry's prefrontal cortex is not learning to control his "panic center", the Amygdala, when he becomes excited or scared. This predicts perhaps lifelong problems with emotional regulation; he may become "a victim of his feelings" later in life, unable to calm down or level his emotional reactions.

Third, the brain will normally produce an entire internal “broadcasting network” from birth to age two, an abundant number of possible dendrite connections far exceeding what we will ever need in life. From age two to puberty, much used connections will be stronger, and not used connections will deteriorate – the brain specializes. In Henry’s case, the lack of early physical stimulation will result in a 20 % brain weight reduction at age two, and the “internal network” base for further development will be of much poorer quality (Struble 1978, Siegle 1999).

And fourth, as Henry leaves babyhood, the low level of brain activity will probably cause a shift into hyperactive, aimless and restless activity because his frontal lobes are not active enough to dominate the rest of the brain. If the lack of stimulation has been very serious, he may have developed “institutional autism”, being very passive and without any wish for contact with others. At twenty months, it is about the last call for changing Henry’s disorganized attachment pattern into a healthier one (Ackerman & Dozier 2005).

TODDLER HENRY

At age three, Henry’s mother gives up, and Henry is placed in a foster care family. His new “family” is doing a great job, but Henry’s behaviour is unfamiliar and confusing to them. Henry has no sense of limits, moral or physical. At one moment he can play with the family’s daughter, age two, the next he will try to strangle her. He creates many fight/ flight situations when the foster parents set limits or make demands. At the dinner table he is sometimes refuses to eat, sometimes eats until he vomits. He does not cry when he falls, he refuses to have physical contact or be cuddled; he takes a long time falling to sleep and often gets up in the middle of the night. His dreams and fantasies are populated with volcanoes, dismembered animals and omnipotent villains. If given an instruction, he is able to repeat it by heart, but immediately forgets it as soon as the adult is out of sight. The foster parents have the notion that he either regards them as enemies (he calls his foster mother “the dragon”), or does not need them at all – he seems indifferent to contact and care. In the kindergarten the other children admire him for his “bravery” (he does not mind jumping from a roof to the ground), and at the same time fear him for his violent temper tantrums. He contacts anybody new in a very irresistible manner, and the foster parents feel misunderstood by their usual environment, who finds him to be a nice, intelligent and charming boy. He quickly makes acquaintances, but no close friendships develop from them. After one year of heroic effort, the foster family gives up when Henry pushes their daughter down the stairs and her arm is broken. Confronted with his actions, Henry’s only comment is: “She stood where I wanted to walk – why did she do that? Should we play again, I’d like that?”.

General aspects: The problem of attachment usually calls for social attention from age two and onwards. It then becomes clear that emotionally and socially the child develops only at a very slow rate, whereas motor and “intelligence” skills may be relatively unaffected (Rygaard 1998). This contradiction in skills often leads to a too optimistic first-meeting view of the child, based on the child’s positive appearance and apparent capacities. However, those who know the child in longer and more intimate relations find out, that the child is imitating the behaviour of others without any understanding of the meaning or intention in that behaviour. Also, that the defence mechanisms in the child are activated by intimate situations that would normally make the child feel safe and loved. These are: fight/flight/freeze patterns, such as vagabonding, attempts to control others – including role reversal in the family, splitting and extremely ambivalent relations – trying to

approach and avoid the adult at the same time. The child will function poorly in socially complex situations (many people, more than one task at a time, new environments or topics), and try to avoid demands or any change in the environment (such as new kinds of food).

HENRY AT SCHOOL

Henry's social worker notices that her hair is turning grey as his case grows heavier. At seven he is placed in another foster home with more experienced foster parents who get regular supervision, and things seem to be all right...

Henry starts in the local countryside school in a positive manner, after a while there are a number of incidents in his disfavour. His teacher is an experienced woman, and her 6 month report is very clear and matter of fact:

"Henry is a very bright and charming little boy, somewhat small for his age. He is able to learn very quickly by heart, but easily forgets what we just talked about. In class, we have had some rather severe conflicts but Henry has found out that I am the leader, and he now obeys somewhat reluctantly. He responds positively to clear and short instructions directed towards him only, he is still unable to attend to a general message to class. In fact he often tries to do what I ask him to do, but he is not able to concentrate and is distracted by what goes on around him, and he is often bored, restless and impatient. He is able to learn quite a lot by copying and repetition, but he has a very concrete way of thinking and does not always grasp the deeper meaning of the issue or text in question. In class he is fairly calm as long as he has my undivided attention, but any new event or unstructured situation turns him upside down, so he still is in conflict outside lessons. He has to be prepared many times each day for what is happening and will happen in the next few minutes. His learning depends on that I break down any subject into little bits which can be learned by heart, one by one. Outside class he now has an "attachment person" who helps him socialize in an acceptable manner."

General aspects: Being more or less intelligent, the resources for learning are often not unfolded in the AD child because of social problems and conflicts overshadowing the learning process. The basic learning problems in AD children all have to do with a failure in the first learning process: perceiving the caregiver as a stable emotional entity. These learning problems (emotional at first, in the cognitive sense later) stem from a lack of the first meaningful exercise in a secure relation:

- concentration (*caregiver responds to contact and prolongs contact*)
- perceiving figure/ background (*caregiver is the emotional "figure"*)
- proportion (*caregiver responds more or less to different events*)
- recognition (*caregiver is the same in varying situations*)
- graduation (*caregiver is empathic and varies stimulation*)
- endure frustration (*caregiver provides safety when situation is uncomfortable*)
- mutual attention (*caregiver is attentive and rewards attention*)
- motivation (*copies caregiver's emotional states, such as joy*)

In other words, object constancy - learned in interaction with the first object/ caregiver – is the prerequisite for a coherent and stable perception of the emotional, physical and social world around us (Blatt 1988).

HENRY THE PERPETUAL TEENAGER

After some rather good years in the foster family Henry enters puberty. He is suddenly caught outside school hours for shoplifting, and the family discovers that this has been going on for a year, including some drug abuse. When confronted his strategy is denial and varying fantastic stories about what happened. He is increasingly angry with his foster parents and refers any problem to them having let him down and not trusting him. In any incident he invariably blames others and is unable to see how he himself influences a situation. There are a number of physical conflicts where the foster parents have to hold him physically in order to avoid his aggression.

His foster mother is beginning to fear him, and while he still seeks comfort like a child he is also looking for weaknesses and feelings of guilt he can awake in them. After one row, he tries to set the house on fire and runs away. He is found some weeks later in a gang of much older youngsters and is brought home by the police. After a calm period the foster parents are called up by a parent from school because Henry has been sexually aggressive towards a girl of eight, and Henry is expelled. At 15 he runs away for good without notice and sometimes surfaces in different court rooms and juvenile institutions.

General aspects: If treated, the child will often have a calm period from the start of school to the start of puberty. Depending on the intensity of the puberty turmoil and the tolerance in the environment, the course of events can be a criminal career (usually fraud, impulsive and aggressive actions and sexual offences due to a low ability to discriminate between appropriate and inappropriate partners), especially if drugs are involved – or if protected and being followed up intensely, a more restricted life with reasonable demands can give a positive outcome. At age 25 the AD child will often demonstrate the emotional maturity level of age 12-14.

HENRY GROWS UP

Even though Henry is no longer criminal at 29, he has a number of problematic behaviours. He often changes jobs, last name and partners and moves around in the country, being restless and easily bored with routine. He does not have a realistic view of himself and he often looks for leader or expert jobs. He is often hired at first glance because he is able to imitate any role in a superficial manner, but also fired within a short time. He likes “getting married” and has produced a numerous offspring and disappointed ex-wives in different cities. If he stays too long with a partner and gets emotionally involved, he sometimes becomes violent and vengeful. At 35 he seems to calm down somewhat: he has a relationship for some years and a regular job.

General aspects: When tested in youth and again at age 40-45, the number of psychopathic traits decreases with age. This is probably due to a “late maturation” of emotional and social function (apart from the mere exhaustion from a very turbulent and exposed life). These traits are:

- Impulsivity

- Superficial and short contacts
- “Charming” behaviour
- Grandiose self-esteem
- Restlessness
- Lies “without reason”
- Lack of guilt and responsibility
- Parasitic lifestyle
- Lack of empathy

(from Robert D. Hare’s Psychopathy Checklist (Hare 1985))

Depending on how and when interventions are made, they may prevent an AD development or reduce the symptoms considerably. Early intervention will of course be most effective.

Unfortunately, the behavioural consequences of deprivation and disorganized attachment only draw attention and concern from age three and onwards, thus producing a lot of practices aimed at treating symptoms that have already become more or less chronic.

POSSIBLE INTERVENTIONS FROM PREGNANCY TO ADULTHOOD

PRE – BIRTH INTERVENTIONS

The first intervention requires a general mapping process in social welfare systems:

Can we point out families whose behaviour makes it likely that the child will develop disorganized/ disoriented attachment?

1. Families with a tradition for criminal behaviour, maltreatment and neglect of children.
2. If substance abuse (alcohol, drugs) also runs in the family, this increases the risk.
3. The mother is not necessarily young, rather she has a destabilized personality (due to her own record of early deprivation/ maltreatment; or to psychosis when the child is in the most basic phase of attachment: age zero to age two). The father is physically or spiritually absent, or violent.
4. Anorectic mothers are at risk of harming the foetus due to malnutrition and problems with attachment after birth.
5. Severely mentally retarded parents.
6. The birth weight of the child is very low/ severe birth complications.

These families/ mothers to be will often already be known in social security and other systems.

Can we direct an intervention program towards these families/ mothers at the onset of pregnancy?

One intervention is to offer mothers-at-risk a special birth preparation program (Lier 1995). In Denmark (5 mio. population) there are currently 48 local programs aimed at mothers-at-risk after many years of focusing on older children.

The general idea of these programs is to give the mother a few stable contact persons, who act as parental figures, helping her cope with daily practical problems, containing and moulding her expectations for motherhood, and help her reduce eventual abuse problems in order to improve the foetal environment. In the study by Lier at Bispebjerg hospital, mothers admitted to such a program produced babies with normal birth weight and a frequency of birth complications not exceeding that of the population in general. Mothers not included demonstrated a reduced birth weight (by 300 grams) and a high frequency of severe complications (50 % of births in sample).

Another program from a small county may be of interest: a health nurse, a psychologist, a social worker, a nursery worker work in a team. When a pregnant at-risk mothers name is referred to the team, she immediately has an informal visit from a team member, who becomes her contact person. She is encouraged to work at a special day care centre for children at risk, where the team is also situated. Here, she is supported by the team who is responsible for her work situation and social welfare benefits; she will “work” in a special day care unit and learn the basics of baby care, she is offered therapy and after birth she will be nursing her own child in the day care centre under staff supervision.

The general idea of this intervention is that the team acts as a “parental substitute”, partly administrating all public aspects (social security, working market, hospital contact, etc.) of the mother’s situation.

The goal of the team is to support the mother and baby until the child is age two (and a healthy attachment process has been started between them). In severe cases, the team helps the mother place the child in a foster family or helps her after mandatory placement in custody of her child. The baby may also be placed in a private day-care where the day-care person has no more than two children, and receives supervision at the day-care centre.

A simpler model is in a county that has a number of specially educated day-care mothers who only work with one or two children at a time. Their job is to be the main attachment figure for the babies, and the biological mother is encouraged to let her baby stay in the day-care for most of the day.

The main idea of all these programs is to provide the mother with a “parental” figure – or a “safe base” - during pregnancy and birth; and to provide practical knowledge and substitute caretakers in the first period of attachment.

INTERVENTIONS FOR MOTHER/ CARETAKER AND BABY RELATIONS

In the intervention programs described, an understanding of healthy neurological development is embedded since a number of studies have demonstrated the close relation between caretaking behaviour and brain development in the baby.

Stimulation of the skin and the vestibular sense (rocking, turning, rolling, etc.) seem to have a major influence on the smooth regulation of the general activity level in the baby brain, and thus both on brain growth and the gradual stabilization of brain function. Consequently, a common problem in AD children is hypo- or hyperactivity. The regulating activity is mediated by the Reticular Activation system, which – in order to function properly in the baby – can be activated only by frequent touch (skin, mouth and tongue) and movement.

A number of caretaker behaviours become important:

1. Frequently carry the baby on the body.
2. Feed the baby in the breastfeeding position on the arm, and spend a lot of time on this. With young fragile babies, a wet nurse can be a good idea.
3. Use baby massage.
4. Exercise eye contact only when the baby is touched at the same time. Touch helps the baby focus and have eye contact.
5. Use a cradle or a hammock for sleeping, never a bed.
6. Let the baby rest on a lambskin or on terry cloth.

Deprived babies quickly adapt to under stimulation and avoid normal levels of stimulation and stay under stimulated. Gradually stimulate the baby for short whiles. Sensory Integration Therapy (Jean Ayres) is designed to teach children how to process rising increasing levels of stimulation.

The Marte Meo program from Holland consists of video feedback sessions for mothers at risk and their babies. Mother and baby are videotaped, and the supervisor gives feedback supporting relevant maternal caretaking behaviour. The general idea of the program is to give only positive feedback whenever the mother displays relevant caretaking behaviour (see: www.martemeo.com).

INTERVENTIONS FOR THE PRE-SCHOOL CHILD

The abovementioned stimulation methods should be put to use even with toddlers and up to age 6, but of course they will be most efficient at an early age.

With AD preschool children, the emotional and social capacities are comparable to those of a much younger child. You should therefore:

1. When planning social and emotional demands and caretaker behaviour, divide the child's age by two or three (to match the developmental age of the child).
2. Do not focus on emotions, reasoning and motivating. Demonstrate the behaviour you want in simple, short sequences and let the child learn them by imitating you instantly (that's how babies learn from their mothers). Don't expect the child to understand the meaning of actions.
7. Keep the child in a "mother/ child" circle wherever you go. That is, the child is followed and helped as closely as a mother would follow her baby. All other social contacts should be planned and supported by the caretaker. AD children are vulnerable to new contacts and large groups.

8. Take responsibility for all negative outcomes of social interaction (wouldn't you do that with a one year old?)

As the reader will remember, our case Henry was placed in a foster care family. Since AD child behaviour is disorganized and a very stressing factor in family life, the following criteria for placement have proven valuable:

1. Foster parents should be experienced, that is, not too young. If they have children the youngest child should be at least 5 years older than the AD child placed in the family.
2. Foster parents should have a stable life without too many new activities and new social contacts. They should be chosen for patience, endurance and stability.
3. Relations to neighbours/ school/ day-care should be positive before placement, since these relations will be stressed by the child's behaviour.
4. Authorities responsible for placement should provide regular supervision.

Mary Dozier of Delaware University has made some very interesting field studies in foster family placement. Some of her most important findings:

- A. If the child is placed in foster care before age 20 months, most children will adapt to the attachment pattern of the foster mother. If placed later, it will probably maintain the attachment pattern acquired from previous caretakers/ parent.
- B. If the foster mother has a secure/ autonomous attachment pattern, the young child placed will adapt to that pattern. If the foster mother has one of the three other attachment patterns (avoidant, ambivalent or disorganized), most children will end up having a disorganized pattern.
- C. Foster parents can learn the behaviours associated with secure/ autonomous attachment.

And other results produced by Femmie Juffer of Leiden University, The Netherlands, studying adopted children (Juffer 2005): if adoptive parents have a number of consultations with an advisor about common attachment problems for one year after adoption, more children will display a secure/ autonomous attachment, compared to a control group of non-advised adoptive parents. The same probably goes for foster parents. It is just as important to attachment outcome to "care for the caretakers" as it is to care for the AD child.

INTERVENTIONS FOR THE SCHOOL-CHILD

The interventions recommended here focus on school work settings, due to the author's experience; that the age five to twelve years is often a reasonably calm period, and that the child often has unused intellectual reserves that are overshadowed by social conflicts. However, the thoughts presented refer to learning processes in general.

To identify the nature of learning problems in AD children, the most relevant cause is the lack of early organized communication patterns between the baby and the caretaker. This prevents the creation of a clear internal representation of the mother, in some theories referred to as "internal working model", in others as "the object relation". In fact, early disorganized interplay probably prevents an ordered experience, not only of the first caretaker, but also of all later persons or

objects, matters and so forth. As mentioned earlier in this text, the subsequent learning problems are caused by poor:

- concentration
- perceiving figure/ background
- proportion
- recognition
- graduation
- endure frustration
- mutual attention
- motivation

These problems appear both in the social relation with the teacher, with peers, and cognitively when working with the understanding of subject matter, such as figures, letters, grammar, ideas and symbols.

Interventions should be designed to support these functions as much as possible, and they should also be the underlying target of exercise in all situations. For example, the duration of concentration can be the teacher's focus in any activity or learning process.

The teacher should regard his or her person more as the "parental object" than as "the teacher", and think in terms of a given number of "parent – child relations" rather than addressing the group of children in general. In other words most of the communication should be between teacher and pupil and not between teacher and group. Only when each child has been in this relation for a year or so can the teacher start activities and instructions on group level.

As a "parental object", the teacher should be aware that he or she only influences the child's behaviour when being present and visible to the child. As soon as the teacher is not present, there will be little or no retention of the teacher's opinions, instructions, etc. In short "you exist only when you are in the presence of the child". For this reason instructions or dialogues about what the child must remember later should be omitted, and only the present tense and the immediate future should be used.

The class room environment should be designed to help the child focus relevantly and not be disturbed, that is, all children face the teacher, and if a child is easily distracted a sound isolating "wall" can separate the children so that a child can only see the teacher.

Concerning the arrangement of learning matter "peeling the onion" is relevant. This means that the teacher initially observes the faculties of the child and only makes demands when he or she is absolutely certain that the child can already perform the task in question. Even then, there will probably be a number of conflicts, because the child perceives the teacher as "the hostile parent" from the original experience of the parent, and this evokes also the aggressive, disorganized or avoidant reactions from this early period. Only when these conflicts have been contained and calmly resolved by the teacher will the child be able to work. It is difficult to be a safe authority without being authoritarian, and this usually takes some years of practice.

What makes teaching with AD children difficult is not only that the internal working model of the child may be imprecise or blurred, but also early adverse experiences will load the child's view of

the teacher with negative and hostile feelings and many defensive mechanisms will blur the communication when the child tries to avoid or disrupt communication. Thus, a calm teaching process in itself is the ultimate goal, what is required is that the teacher learns to be a “container for negative projections” (that is to know that the child’s problems stem from early life and are not really statements to be taken personal) and also learns to be a good “rodeo cowboy” (that is, the child will often try to create a secure base by role reversal or by taking control of the environment).

It is equally important to understand that the teacher’s efforts to explain or convey the deeper meaning or essence of a subject can be an insurmountable challenge for the child. Instead, the logics of a task should be translated into a set of behaviours and rituals which the child can simply imitate – the “why” should always be replaced with the “how” demonstration (Rygaard 2006).

INTERVENTIONS IN YOUTH

This presentation has focused on early childhood and school years, and youth has been described in many other places. There are so many intervention programs that it is impossible to give a fair description. In general, programs based on cognitive/ behavioural therapies are popular in these years. Without any evidence, it is my experience that these programs work well with many “light” youth criminals, but as with all other methods they have only a temporary effect on youngsters with many psychopathic traits. In these cases, an intense permanent follow up and control procedure is necessary. In a Danish facility for psychopathic criminals (Herstedvester), a permanent follow up is combined with re- incarceration without time limit if a parole is violated. The next parole then depends solely on the estimate of prison staff. This method seems to have some effect in the most difficult cases.

CONCLUSIONS

What has amazed me mostly in working with AD children is the fact that the causes seem to operate from pregnancy to age two, while treatment usually starts when the social symptoms start to become increasingly manifest: from age 2-3 and onwards. Due to the late start of treatment, the prognosis is not very positive. This calls for the development of intervention programmes early in life.

Milieu therapy, in my experience, is a useful method in providing the secure base to extremely insecure and aggressive children. The two most important goals in this are to monitor the environment at all times (which can usually be obtained at least until puberty even in severe cases), and to provide supervision for those who work with the children in order to avoid regression under the pressure of the child’s behaviour problems.

*Source: Rygaard, N.P.: “Severe Attachment Disorder in Childhood – a Guide to Practical Therapy” Springer Wien/ New York 2006. www.springer.com
“L’Enfant Abandonné – Guide pour le Traitement de Troubles de l’attachement”,
De Boeck Université 2006. www.deboeck.com*

See/ Voir: www.attachment-disorder.net, mail author: npr@erhvervspsykologerne.dk

LITERATURE

For the reader's inspiration for further reading, the literature list from "Severe Attachment Disorder in Childhood" is presented in its full length. The studies referred to in the above text are included here. The references in this paper to Ackerman (2005) and Juffer (2005) are not included in the book.

- Ackerman, J. P., & Dozier, M. (2005). The influence of foster parent investment on children's representations of self and attachment figures. *Applied Developmental Psychology, 26*, 507-520.
- Ainsworth M (1969): Object Relations, dependency and attachment: A theoretical review of the infant-mother relationship. *Child Development, 40*: 969 – 1025
- Ainsworth M, Blehar, M, Waters I, Wall S (1978): *Patterns of Attachment – a Psychological Study of the Strange Situation*. Lawrence Erlbaum Associates, Hillsdale
- Amtsraadsforeningen (2000) *The public school special need classes 1985-1999*. Amtsradsforeningen, Denmark
- Ayres J (1979) *Sensory integration and the child*. Los Angeles, CA
- Barnard KE & Brazelton TB (1990) *Touch – the foundation of experience*. International Universities Press, Madison, Conn.
- Berman AJ, Berman D. & Prescott JW (1974) The Effect of cerebellar lesions on emotional behavior in the rhesus monkey. In: Cooper, I.S., Riklon, M.V. & Snider, R.S. (eds) *The cerebellum, epilepsy and behavior*. Plenum, NY
- Bion WR (2000) *Experiences in groups and other papers*. Routledge, London.
- Blatt SJ (1988) A cognitive morphology of psychopathology. In: plenary lecture at the ISSPD congress on personality disorders, Rigshospitalet, Copenhagen
- Blatt SJ (1988) Interpersonal relatedness and self - definition. IN: Singer JL (ed) *Defense mechanism and personality style*. University of Chicago Press, Chicago
- Bowlby J (1969) *Attachment and loss, vol. 1. Attachment*. Hogarth Press, London. Basic Books, New York; Penguin, Harmondsworth
- Bowlby J (1973): *Attachment and Loss, vol. 2. Separation: anxiety & anger*. Hogarth Press, London. Basic Books, New York; Penguin, Harmondsworth
- Bowlby J. (1988) *A secure base: Clinical applications of attachment theory*. Routledge, London
- Briggs F & Hawkins RMF (1996) A comparison of the childhood experiences of convicted male child molesters and men who were sexually abused in childhood and claimed to be non - offenders. In: *Child Abuse and Neglect, 20*: 221 -233.
- Bryan GK and Riesen AH (1989) Deprived somatosensory-motor experience in stump-tailed monkey neocortex: dendritic spine density and dendritic branching of layer III B pyramidal cells. *J Comp Neurol 286*: 208-217
- Coleman M (1971) Platelet serotonin in disturbed monkeys and children. *Clinical Proceedings of the Children's Hospital. 27 (7)*: 187-194
- Deutch AY & Roth RH (1990) The determinants of stress induced activation of the prefrontal cortical dopamine system. *Progr Brain Res 85*: 367 – 404
- Dokecki PR (1973) When the bough breaks...what will happen to baby. Review of: *Rock-a-bye Baby*. Time Life Films (Lothar Woff, Ex. Prod.) *Contemporary Psychology 18*:64
- Doren DM (1987) *Understanding and treating the psychopath*. Wiley, New York
- Essman WB (1971) Neurochemical changes associated with isolation and environmental stimulation. *Biological Psychiatry, 3*: 141

Field T, Hernandez-Reif M, Seligman S, Krasnegor J, Sunshine W, Ricas-Chacon R, Schanberg S, Kuhn C (1997) Juvenile rheumatoid arthritis: benefits from massage therapy. *J Pediatr Psychol*, 22: 607-617

Field T, Hernandez-Reif M, Diego M, Feijo L, Yanexy V & Gil K (2004) Massage therapy by parents improves early growth and development. *Infant Behavior and Development*, 27: 435 – 442

Finkelhor D, et al (1990) Sexual abuse in a national survey of adult men and women: Prevalence, characteristics, and risk factors. *Child Abuse and Neglect*, 14: 19 - 28

Fleming J et al (1997) The long term impact of sexual abuse in Australian women. *Child Abuse and Neglect* 21, (1): 49 - 58

Fonagy P Pathological attachments and therapeutic action. Paper to the Developmental and Psychoanalytic Discussion Group, American Psychoanalytic Association Meeting, 13 May 1999, Washington, DC

Gallahue DL (1997) Understanding motor development: infants, children, adolescents, adults, 5th edn. WCB Brown & Benchmark, Madison

Green A et al (1981) Neurological impairment in maltreated children. *Child Abuse and Neglect* 5: 129 -134

Hansen N (1977) Cerebro-organic Pathogenesis in 110 Children followed up subsequent to Admission to a Child Psychiatric Ward. *Acta Psychiatrica Scandinavica* 46: 399-412

Hare RD (1985) The psychopathy checklist. University of British Columbia, Vancouver

Hare RD (1988) Psychopathy and language. In: Moffit TE & Mednick SA (eds.): *Biological contributions to crime causation*. Nijhoff, Dordrecht

Hare RD (1993) *Without conscience: The disturbing world of the psychopaths among us*. Simon & Schuster, New York, NY (reissued in 1999 by Guilford Press)

Hare RD, Forth AE, & Strachan, KE (1992) Psychopathy and crime across the lifespan. In Peters R, McMahon R, & Quinsey V (eds) *Aggression and violence across the lifespan*. Sage, Newbury Park, CA, 285-300

Hart SD, Cox DN & Hare RD (1995) The Hare psychopathy checklist: Screening version.: Multi-Health Systems, Toronto, ON

Heath RG (1972) Electroencephalographic studies in isolation raised monkeys with behavioral impairment. *Dis Nervous Systems*, 33: 157-163

Heath RG (1975): Maternal-social deprivation and abnormal brain development: Disorders of emotional and social behavior. In: Prescott JW, Read MS, Coursin DB (eds) *Brain function and malnutrition. Neuropsychological methods of assessment*, Wiley, New York

Higley JD, Suomi SJ, Linnoila M (1990) Parallels in aggression and serotonin: Consideration of development, rearing history, and sex differences. In: van Praag H, Plutchik R, Apter A (eds) *Violence and suicidality: Perspectives in clinical and psychobiological research*. Brunner/ Mazel, New York

Hoksbergen RAC, Rijk JK(2004) Effects of deprivation. An example: Children adopted from Romania. Utrecht University, Department Adoption/Non-Genetic Parenthood, Heidelberglaan 2, 3584 CS Utrecht, The Netherlands

James B (1994) *Handbook for treatment of attachment - trauma problems in children*. The Free Press, New York

Juffer F Bakermans-Kranenburg, MJ., & Van IJzendoorn MH (2005) The importance of parenting in the development of disorganized attachment: Evidence from a preventive intervention study in adoptive families”, *Journal of Child Psychology and Psychiatry* 46, 263-274.

Kernberg O (1996) *Personality development and psychopathology*.
Video recording: Lecture at The Psychiatric Dpt. Of Roskilde Hospital, Denmark.

van der Kolk B & Greenberg MS (1987) The psychobiology of the trauma response: Hyperarousal, constriction, and addiction to traumatic re-exposure. In: van der Kolk B (ed) Psychological trauma. American Psychiatric Press, Washington, DC, pp 63-87

van der Kolk B (1994) Trauma and the development of borderline personality disorder. *Psychiat Clin North Am* 17, (4): 715 - 730

Kruuse E (1984) Skoleforløbet for børn med lav fødselsvægt (Learning disabilities and education in children with low birth weight), Skolepsykologisk Forlag, Copenhagen

Laudenslager ML, Reits M, Harbeck R (1982) Suppressed immune response in infant monkeys associated with maternal separation. *Behav Neural Biol* 36: 40-48

Leth I (1999) Seksuelle overgreb mod børn, karakteristika og psykiske skadevirkninger. (Sexual assault towards children, characteristics and psychological damage). In: *Psykkiske sygdomme og problemer hos børn og unge (Psychiatric Disease and Problems in Children and Juveniles)*, Psykiatrifondens Forlag, Copenhagen 1999, pp 205-218

Lier M, Gammeltoft M, Knudsen IJ (1995) Early mother-child relationship: The Copenhagen model of early preventive intervention towards mother-infant relationship disturbances. *Arc Medical Res*, 54 (Suppl 1): 15-23

Lisak D (1994) The psychological impact of sexual abuse: Content analysis of interviews with male survivors. *J Traumatic Stress*, 7: 507 - 523

Madsen SA (1996) Baand, der brister – baand, der knyttes. (Bonds that break, bonds that tie). Reitzel, Copenhagen

Main M (1996) Introduction to the special section on attachment and psychopathology: 2. Overview of the field of attachment. *J Consult Clin Psychol* 64, (2): 237 – 243

Mason WA (1968) Early social deprivation in the non-human primates: Implications for human behavior. In: Glass DE Ed *Environmental influences*. The Rockefeller University Press/ Russell Sage Foundation, New York

Mason WA, Berkson G (1975) Effects of maternal mobility on the development of rocking and other behaviors in rhesus monkeys: A study with artificial mothers. *Dev Psychobiol* 8: 197-221

McCord W (1982) *The Psychopath and milieu therapy: A longitudinal study*. Academic Press, New York

Meloy JR (1988) *The psychopathic mind*. Jason Aronson, London

Melzack R, Burns SK (1965) Neurophysiological effects of early sensory restriction. *Exp Neurol* 13: 163-175

Montagu A (1986) *Touching - the human significance of the skin*. Harper and Row, New York

Neal M (1967) Vestibular stimulation and developmental behavior in the small premature infant. *Nursing Research Report* 3: 1-4

NICHD (National Institute of Child Health and Human Development) (2003) Child Care linked to assertive, noncompliant and aggressive behaviors – vast majority of children within normal range. [Http://www.nichd.nih.gov/new/releases/child_care.cfm](http://www.nichd.nih.gov/new/releases/child_care.cfm).

Piaget J (1936/ 1953) *Origins of intelligence in the child*, Routledge & Kegan Paul, London

Prescott JW (1971) Early somatosensory deprivation as an ontogenetic process in the abnormal development of the brain and behavior. In Goldsmith IE, Moor-Jankowski J (eds) *Medical primatology 1970*. Karger, Basel New York

Prescott JW (1975) Body pleasure and the origins of violence. *The Bulletin of The Atomic Scientists*, November 1975, pp 10-20

Prescott JW (1977) Phylogenetic and ontogenetic aspects of human affectional development. In: Gemme R, Wheeler CC (eds) *Progress in Sexology. Proceedings of the 1976 International Congress of Sexology*. Plenum Press, New York.

Prescott JW (1979) Deprivation of physical affection as a primary process in the development of physical violence. In: Gil DG (ed) *Child Abuse and violence* AMS Press, New York, pp 66-137

Prescott, J.W. (1980). Somatosensory affectional deprivation (SAD) theory of drug and alcohol use. In: Lettieri DJ, Sayers M, Wallenstien Pearson H (eds) *Theories on drug abuse: Selected contemporary perspectives* (NIDA Research Monograph 30, March 1980). National Institute on Drug Abuse, Department of Health and Human Services. Rockville, MD.

Prescott JW (1990) Affectional bonding for the prevention of violent behaviors: Neurobiological, psychological and religious/spiritual determinants. In: Hertzberg LJ, et al (eds) *Violent behavior, vol I: Assessment and intervention*. PMA Publishing, NY pp 110-142

Reid WH, Dorr D, Walker J & Bonner J (1986) *Unmasking the psychopath: Antisocial personality disorder and related syndromes*. Norton, New York

Reite M & Capitanio JP (1985) On the nature of social separation and social attachment. In: Reite M, Field T (eds) *The psychobiology of attachment*. Academic Press, New York

Riordan D, Appleby L, Faragher B (1999) Mother-infant interaction in post-partum women with schizophrenia and affective disorders. *Psychological Medicine* 29, 991-995

Rosenzweig MR (1971) Effects of environment on development of brain and of behavior. In: Tobach E, Aronson LR & Shaw E (eds). *The biopsychology of development*. Academic Press, New York, 303-342

Rutter M (1995) Relationships between mental disorders in childhood and adulthood. *Acta Psychiatrica Scandinavica* 91: 73-85

Rygaard NP (1998) Psychopathic children: Indicators of organic dysfunction. In: Millon T, et al (eds) *Psychopathy. Antisocial, criminal and violent behavior*. Guilford Press, New York, pp 247-259

Rygaard NP (2004): A systemic approach to cross-scientific terms in the diagnostic Babylon. (unpublished)

Salk L, Lipsitt LP, Sturner WQ, Reilly BM & Levat RH (1985) Relationship of maternal and perinatal conditions to eventual adolescent suicide. *Lancet*, 1985 Mar 16;1(8429):624-627

Schalling, D. (1988): *Personality Self-Report Scales and Biological Markers for Aggressivity and Lack of Behavioral Constraints*. Plenary Lecture at: First Congress of the International Society for the Study of Personality Disorders. Copenhagen

Schore AN (1994) *Affect regulation and the origin of the self: The neurobiology of emotional development*. Lawrence Erlbaum, Hillsdale NJ

Schore AN (2002) Dysregulation of the right brain: A fundamental mechanism of traumatic attachment and psychopathogenesis of posttraumatic stress disorder. *Austr NZ J Psychiat* 36: 9-30

Schulsinger F (1972) Nogle undersøgelser til belysning af sammenhængen mellem arv og miljø i psykiatrien. (Some studies concerning the link between hereditary and milieu factors in psychiatry). Foreningen af danske lægestuderende, Copenhagen

Schutz WC (1958) *FIRO: A three-dimensional theory of interpersonal behavior*. Holt & Rhinehart, New York (Re-issued in 1966 as *The Interpersonal Underworld* by Science and Behavior Books, Palo Alto, CA)

Shaver PR, Cassidy J (eds) (1999): *Handbook of Attachment – theory, research, and clinical applications*. Guilford Press, New York

Siegel DJ (1999): *The developing mind – toward a neurobiology of interpersonal experience*. Guilford Press, New York

Struble RG, Riesen AH (1978) Changes in cortical dendritic branching subsequent to partial social isolation in stump-tail monkeys. *Dev Psychobiol*, 11(5): 479-486

Tassinari CA (1968) Suppression of focal spikes by somatosensory stimuli. *Electroenceph Clin Neurophysiol* 25: 574-578.

Teicher MH. et al (1987) Preliminary evidence for abnormal cortical development in physically and sexually abused children using EEG coherence and MRI. In: van der Kolk B. & Greenberg MS (1987): *The Psychobiology of the trauma response: Hyperarousal, constriction and addiction to traumatic re-exposure*. American Psychiatric Press, Washington, D.C, pp 63-87

Teicher MH et al (1994) Early abuse, limbic system dysfunction, and borderline personality disorder. In: Silk K (ed): *Biological and neurobiological studies of borderline personality disorder*. American Psychiatric Association Press, Washington, DC, pp 177-207

Turnbull CM (1987) *The mountain people*. Simon & Schuster, New York

Vanggaard T (1968) Neurose og Psykopati (Neurosis and psychopathy). In: *Nordisk Psykiatrisk Tidsskrift*, 4: 22-30, Copenhagen.

Viding ER, Blair JR, Moffitt TR, Plomin R (2005): Evidence for substantial genetic risk for psychopathy in 7-Year olds. *J Child Psychol and Psychiat* 46:6, pp 592 – 597

White W (1959) Motivation reconsidered - the concept of competence. In: *Psych Review*, vol. 66, pp 297-333

Williams LM (1994) Recall of childhood trauma: A prospective study of women's memories of child sexual abuse. In: *J Consult Clin Psychol* 62, pp 1167 - 1176

Williamson SE, Harpur TJ, Hare RD (1991). Abnormal processing of affective words by psychopaths. *Psychophysiology* 28 (3): 260-273

Zachau-Christiansen B (1975) *Babies: Human development during the first year*. Wiley, Chichester

